

Improving manure recycling in agriculture – an important step towards closing the phosphorus cycle

Phosphorus is a finite resource; efficient usage and recycling are essential. Recycling human sewage can help close the phosphorus cycle, but only partially meets circular economy goals. Phosphorus recycling efforts must not neglect manure.

All life depends on phosphorus. But it is a finite resource for which there is no substitute. The inefficient and wasteful use of phosphorus in the agri-food system has implications for sustainability and also contributes to eutrophication of lakes, streams, and coastal areas. The responsible and sustainable use of phosphorus requires better use and re-use in all sectors of society, especially in agriculture.

The EU Circular Economy Action Plan has generated interest in recycling phosphorus contained in human sewage in agriculture. This effort is an important step in closing the phosphorus cycle. However, there is greater potential to improve the efficiency of manure used as crop fertiliser. In the Baltic Sea catchment, the amount of phosphorus in manure is more than 3 times greater

than in human sewage. By using manure more efficiently, mineral fertiliser imports could be reduced by 0,11 - 0,17 million tons, compared to about 0,036 million tons by using sewage sludge. This would reduce phosphorus surpluses and the risk of leakage to inland waters and the Baltic Sea.

Challenge to meet Baltic Sea Action Plan reduction targets

[Progress has been made](#) in reducing nutrient inputs to the Baltic Sea. Since 1995, nitrogen inputs have decreased by 22% (250 000 tons) and phosphorus by 33% (15 000 tons).

To achieve the goals of the Baltic Sea Action Plan, [further nutrient reductions are needed](#). Compared to average inputs for the 2012-2014 period, nitrogen inputs must decrease by 12% (122 000 tons) and phosphorus by 55% (14 000 tons) to reach maximum allowable inputs.

Nutrients that cycle in the agri-food system contribute to eutrophication by, for example, leakage of fertiliser and manure, which have been applied to crop land, and discharge of sewage effluent to water bodies.

Past inefficient use has led to the accumulation on land in the past 50 years, countries around the Baltic Sea have applied about 35 million tons of mineral phosphorus fertilisers to agricultural soils.

Decades of inefficient use of phosphorus in crop production has led to the accumulation of huge amounts of nutrients in soils. Earlier, farmers were advised to apply large amounts of



mineral fertiliser and manure to build up soil phosphorus levels. Soils in several regions now contain so much phosphorus that further fertilization does not increase harvest.

For the Baltic Sea catchment as a whole, the magnitude of soil phosphorus surplus represents about two decades of crop nutrient needs ¹. Consequently, here is potential to decrease application rates of mineral fertilisers and manure in areas with substantial phosphorus surpluses.

This accumulation, or so-called surplus, increases the risk of losses to lakes, streams, and the Baltic Sea. And the accumulated surplus keeps growing.

The buildup of phosphorus on land can be slowed by reducing imports of feed and mineral fertilisers through more effective recycling of nutrients in manure and human sewage and by avoiding over-fertilisation.

Phosphorus use efficiency varies greatly around the Baltic Sea ²

Tons P	Outputs	Inputs		PUE*	Human excreta
	Crop harvest	Fertiliser	Livestock excreta		
Belarus	36 400	69 400	46 300	0,3	5 400
Denmark	45 400	11 200	55 200	0,7	7 200
Estonia	6 400	3 000	3 800	0,9	1 400
Finland	21 600	12 200	15 600	0,8	6 600
Germany	55 000	19 200	34 100	1,0	6 500
Latvia	12 700	7 100	5 900	1,0	2 300
Lithuania	25 300	14 400	12 200	0,9	4 100
Poland	165 100	170 300	154 000	0,5	45 000
Russia	4 500	2 600	16 800	0,2	13 200
Sweden	34 000	10 500	22 700	1,0	11 800
Total	406 400	319 900	366 600	0,6	103 500

*In countries with low phosphorus use efficiency, there is the opportunity to reduce imports of imported mineral fertilisers by avoiding over-fertilization and by substitution with animal manure (excreta). Phosphorus in human excreta can also be further used better recycled in agriculture, further reducing the region's dependence on fertiliser imports. *PUE is phosphorus use efficiency in crop production, the ratio of crop harvest (outputs) to the sum of fertiliser and livestock excreta (inputs)*

Recycling could reduce mineral fertiliser imports

Agricultural systems will never be perfectly efficient because of unavoidable nutrient losses. But in the Baltic Sea region, there is room for improvement. Each year, the 23 million pigs, 16 million cows, and 244 million chickens in the region produce manure that contains over 0,35 million tons of phosphorus². Much of the phosphorus in the manure originates from imported animal feed. Even though over 70% of crops grown in the Baltic Sea catchment are fed to livestock, this is insufficient to meet the animals' nutrient requirements. As a result, two-thirds of the phosphorus needed by livestock is imported, mostly as soy from South America^{2,3}.

A recent study shows that the average phosphorus use efficiency in crop production in the Baltic Sea region is only about 60%, but it varies greatly between countries. If phosphorus use efficiency increased to 70 - 90%, imports of mineral phosphorus fertilisers could be reduced by 0,1 to 0,2 million tons⁴. Specialisation and spatial separation of crop and livestock production systems can cause excessive phosphorus application in animal-dense regions that have large amounts of manure in relation to crop area⁵. There is potential to use manure more efficiently by redistributing "excess" manure nutrients from regions specialising in livestock production to regions specialising in crop production and, ultimately, reduce mineral fertiliser imports.

Recycling sewage sludge could reduce fertiliser imports

About 75% of people in the Baltic Sea region are connected to centralised sewage systems, while the remainder are connected to non-centralised systems, such as septic- and cesspit-type systems. Centralised sewage treatment practices have improved substantially in the past few decades; however, phosphorus removal efficiencies vary greatly, between 63% (Latvia) and 97% (Finland, Germany, and Sweden)⁶.

There is opportunity to recycle phosphorus from sewage sludge that is collected in centralised systems. Today, only about one third of the sewage sludge in the region, which contains about

0,027 million tons of phosphorus, is used in agriculture, but practices differ among countries in the region⁷. The remaining sludge, corresponding to 0,036 million tons of phosphorus, could be used in agriculture and reduce fertiliser imports⁸.

Phosphorus that is not removed from sewage as sludge in centralized sewage systems is discharged to surface waters in effluent. Thus, improving sewage treatment capabilities is an effective way to prevent phosphorus from entering lakes and rivers that drain to the Baltic Sea. Currently, an average of 16% of phosphorus in human excreta (0,011 million tons) becomes effluent⁷.

Phosphorus is a finite resource

Mineral phosphorus fertilisers derive from phosphate rock, which is a finite resource, like fossil fuels. Unlike fossil fuels, however, there is no substitute. Today, 85% of phosphate rock reserves are controlled by three countries: Morocco, China, and Algeria⁹. The European Commission has added it to the list of 20 Critical Raw Materials for which supply security is at risk and economic importance is high¹⁰.

In fact, society currently operates outside the planetary boundaries of the phosphorus cycle. This is not sustainable^{11,12}. Therefore, recycling existing phosphorus within agriculture better, and reducing the overall import of new phosphorus to the Baltic Sea region, not only mitigates eutrophication but also leads to a more sustainable use of a finite resource.

RECOMMENDATIONS

There is potential to make progress towards the goals of the Baltic Sea Action Plan by increasing the recycling of phosphorus in agriculture and reducing over-fertilisation.

- National actions should not neglect manure.
- Increase the use of locally produced feed in animal husbandry in order to reduce imports of phosphorus to the Baltic Sea region.
- Facilitate the trade of recycled fertilisers by, for example, supporting the proposed EU fertiliser trade legislation.
- Improve the use efficiency of manure use in agriculture by, for example, setting legal limits for fertilisation of soils and supporting farm extension on nutrient management, including soil nutrient mapping.

Scientific sources and notes

- 1 Rowe, H. *et al.* Integrating legacy soil phosphorus into sustainable nutrient management strategies for future food, bioenergy and water security. *Nutrient Cycling in Agroecosystems*, doi:10.1007/s10705-015-9726-1 (2015).
- 2 Hong, B. *et al.* Advances in NANI and NAPI accounting for the Baltic drainage basin: spatial and temporal trends and relationships to watershed TN and TP fluxes. *Biogeochemistry*, doi:10.1007/s10533-017-0330-0 (2017).
- 3 FAOSTAT. Food and Agriculture Organization of the United Nations, FAOSTAT Database Collections. <http://faostat.fao.org/>. (2016).
- 4 Baltic Eye. Estimate made using data from Hong et al. (2017).
- 5 Nesme, T., Senthilkumar, K., Mollier, A. & Pellerin, S. Effects of crop and livestock segregation on phosphorus resource use: A systematic, regional analysis. *European Journal of Agronomy* **71**, 88-95, doi:10.1016/j.eja.2015.08.001 (2015).
- 6 Hautakangas, S., Ollikainen, M., Aarnos, K. & Rantanen, P. Nutrient abatement potential and abatement costs of waste water treatment plants in the Baltic Sea region. *Ambio* **43**, 352-360, doi:10.1007/s13280-013-0435-1 (2014).
- 7 Eurostat. Sewage sludge production and disposal (env_ww_spd). http://appsso.eurostat.ec.europa.eu/nui/show.do?dataset=env_ww_spd&lang=en. Accessed 6 May 2015. (2016).
- 8 Baltic Eye. Phosphorus contained in sludge generated in Russia and Belarus was included with other/unknown because of lack of data.
- 9 Elser, J. J. & Bennett, E. M. A broken biogeochemical cycle. *Nature* **478** (2011).
- 10 EC. The European Critical Raw Materials Review Memo. European Commission (EC), Brussels, Belgium ([http://europa.eu/rapid/press-release MEMO-14-377_en.htm](http://europa.eu/rapid/press-release_MEMO-14-377_en.htm)). (2014).
- 11 Carpenter, S. R. & Bennett, E. M. Reconsideration of the planetary boundary for phosphorus. *Environmental Research Letters* **6**, 014009, doi:10.1088/1748-9326/6/1/014009 (2011).
- 12 Steffen, W. *et al.* Planetary boundaries: Guiding human development on a changing planet. *Science* **347** (2015).

